

**Regulating khat? Day Conference and Panel Debate
Middlesex University, London, 4 March 2011**

***Assessing khat-related death, with special
reference to the UK situation***

John M. Corkery, BA Hons, MSc, MPhil
Senior Research Fellow in Drug Epidemiology, International
Centre for Drug Policy, St George's University of London, UK

Speaker's experience

Currently Programme Manager for the National Programme on substance Abuse Deaths & VSA Mortality Project

UK Focal Point expert on Drug-Related Mortality statistics (since 2000)

UK Focal Point advisor on Availability and Offending indicators

Formerly responsible for the production of Home Office Statistics on notified addicts, drug seizures and drug offences

Give lectures on range of drug statistics, control & epidemiology

(Co)-author of peer-reviewed journal articles, co-editor of 3 international drug monographs, etc

WHO Temporary Advisor to EMRO October 2007 at 6th RAPID meeting in Cairo, presenting on public health issues relating to khat

Overview

Introduction

What is a “khat-related” death?

National Programme on Substance Abuse Deaths

Method and data sources for UK deaths

Characteristics of UK khat-related deaths

Discussion

Comparison with other substances

Acknowledgements

References

Contact details

Introduction

During the past 20 years or so, more has become known about the properties of khat, its pharmacology, physiological and psychological effects on humans. Khat consumption has adverse health consequences including myocardial infarction, liver failure, depression, psychoses, and dependence. However, its reputation of social and recreational use in traditional contexts has hindered the dissemination of knowledge about its detrimental effects in terms of mortality.

This paper focuses on this particular deficit, briefly outlining the types of mortality associated with the trade and use of khat. With khat being increasingly brought under domestic regulation in many countries, and its legal status being considered in others, it is important that both these dimensions need to be understood.

What is a “khat-related” death ?

Prior to our research paper, there were no known published reviews of or statistics on khat-related mortality - an important gap in the knowledge-base.

A literature search of all relevant databases as well as the Internet to identify relevant reports and information on ‘khat-related’ mortality was undertaken. Need media reports for deaths associated with khat trade and consumption; not in academic databases.

This process identified a range of deaths that can be directly and indirectly associated with khat, some of which can overlap.

To help understand the dimensions of the phenomenon it is necessary to derive themes or ‘motifs’ from the results (Table 1).

Table 1: Taxonomy of khat-related deaths

Directness	Type of association	Mechanism	Example
Trade-related			
	Cultivation/production	Poisoning	Fertiliser/Pesticide not washed before consumption
		Disputes between actors	Disagreement over wages => homicide
	Transportation	Fatigue	Long hours and driving at high speed => loss of concentration (can be in association with khat use to keep awake) leading to accidents
		Loss of concentration	Distraction whilst preparing khat for chewing whilst driving => accidents
		External factors	Bad weather conditions/mechanical failure => plane crashes
	Distribution/marketing/wholesale	Disputes between actors	Fighting over 'turf' => violence and homicide
		Disputes between actors	Refusal to do business => violence and homicide
	Retail	Disputes over price	Homicide

Taxonomy of khat-related deaths

Directness	Type of association	Mechanism	Example
Consumption-related (medical)			
	Physiological	Mechanical	Choking on leaves/twigs or airway obstruction => asphyxia or cardiac arrest
		Toxicity	Myocardial infarction => fatal heart attack
			Liver failure
	Neurological	Lack of motor co-ordination, shaking	Reduces control => accidents
		Eye-sight problems	Impairs sight and focus => accidents
	Psychopathological	Causing and/or exacerbating psychosis and/or depression	Suicide and/or homicide, accidental overdose
		Impaired judgement/assessment of risk => accidents	Road traffic accident
			Fall from height
	Other medical contributory factors	Used with other psychoactive substances and/or positive toxicology	Role may not be clear

Taxonomy of khat-related deaths

The contribution of khat to deaths is varied; and in some cases can be difficult to categorise.

Deaths associated with trade, marketing, wholesale and retail, can equally occur with any other illicit substance.

Their inclusion here is deliberate: they are part of the wider pattern of mortality associated with the consumption of khat.

The association of khat with psychiatric disorders or psychopathological factors is evident in some cases, contributing to suicide and even homicide.

The physiological effects of khat consumption are quite clear in a number of cases.

National Programme on Substance Abuse Deaths (np-SAD)

np-SAD gets information from coroners and other sources across the UK on a voluntary basis on drug-related deaths and deaths of addicts.

Since 1997 details of 23,000 deaths received.

The average annual response rate is up to 95% (Ghodse et al., 2009).

Case definition - at least one of the following: (a) presence of one or more psychoactive substances directly implicated in death; (b) history of dependence or abuse of drugs; (c) presence of controlled drugs at post-mortem.

Method and data sources for UK deaths

We defined deaths related to khat as the inclusion of the words 'qat', 'khat', cathine, cathinone, norephedrine or (nor)pseudoephedrine in the cause of death, post-mortem drugs, verdict, and incident description sections of the coroner's report.

The presence of such compounds in the toxicology results was used in conjunction with other information to define a khat-related death, since norpseudoephedrine is also a metabolite of pseudoephedrine which is a precursor for methylamphetamine and can therefore be derived from a substance that is not cathinone.

Method and data sources for UK deaths

Case identification - quasi 'snow-ball' cluster sampling approach.

5 notified as part of the normal surveillance programme. Tox investigations for 2 of these done by SGUL Forensic Toxicology Service. Examination of their records revealed 3 more deaths.

Members of London Toxicology Group asked if they knew any cases => information on one case.

6 were identifiable through press reports; info got from coroners.

Information extracted using np-SAD data form. Details of individual's background, medical & psychiatric history, PM & tox reports obtained/scrutinised at coroner's office by author.

Enquiries were also made of UK General Register Offices & SCDEA to ascertain any other cases – none identified.

Characteristics of UK cases – paper in *Drugs – education, prevention and policy*

Drugs: education, prevention and policy, Early Online: 1–18
Copyright © 2010 Informa UK Ltd.
ISSN: 0968-7637 print/1465-3370 online
DOI: 10.3109/09687637.2010.504200

informa
healthcare

ORIGINAL RESEARCH

'Bundle of fun' or 'bunch of problems'? Case series of khat-related deaths in the UK

John M. Corkery¹, Fabrizio Schifano², Adenekan Oyefeso¹, A. Hamid Ghodse¹, Thomy Tonia³, Vinesha Naidoo¹, & Jennifer Button⁴

¹National Programme for Substance Abuse Deaths (np-SAD), International Centre for Drug Policy, St George's, University of London, 6th Floor, Hunter Wing, Cranmer Terrace, London, SW17 0RE, UK,

²School of Pharmacy, University of Hertfordshire, Hatfield, Hertfordshire, AL10 9AB, UK, ³Institute of Social and Preventive Medicine, University of Bern, Finkenhubelweg 11, Bern, CH 3012, Switzerland, and

⁴Forensic Toxicology Service, Division of Cardiac and Vascular Sciences, St George's, University of London, Ground Floor, Jenner Wing, Cranmer Terrace, London, SW17 0RE, UK

Twenty million people worldwide use khat (*Catha edulis*). Previously confined to Eastern Africa and Arabia, consumption is spreading to other regions. Chewing khat leaves releases the stimulants cathinone and cathine. Khat consumption has adverse health consequences including myocardial infarction, liver failure, depression, psychoses and dependence. Literature regarding khat-related mortality is scant: only one death (in 1945) due to

psychological effects of khat, together with the risks for mortality associated with its use.

INTRODUCTION

Khat (*Catha edulis*) is a shrub that grows in eastern Africa and southern Arabia. Its leaves and twigs are

altheare.com by \${individualUser.displayName} on 09/11/10
personal use only.



Table 2: Main characteristics of 14 khat-related deaths, UK, 2004-9

<i>Characteristics</i>	<i>Frequencies</i>
Year of death	2004 = 2; 2005 = 3; 2006 = 4; 2007 = 0; 2008 = 4; 2009 = 1
Geographical area	East London = 1; West London = 3; North London = 7; Hampshire = 1, Avon = 1, Cardiff = 1
Gender	All male
Marital status	Married = 3; with partner = 1; divorced = 1; separated = 1; single = 2 not known = 6
Age at death (years)	Mean = 36, range = 19 to 47
Ethnicity/Nationality	Somali = 12; Eritrean = 1; Polish = 1
Length of UK residence (years)	3 = 1; 5 = 1; 11 = 2; 15 = 2; Not known = 8
Occupation	Unemployed = 6; manual employed = 4; student = 2; invalidity benefit = 2
Living arrangements	Alone = 3; partner & children = 3; self & children = 1; with partner = 1; with sibling = 1; with cousin = 1; with friends = 1; psychiatric in-patient = 1; not known = 2
Significant medical history	No = 7; yes = 3; not known = 4
Known psychiatric history	Yes = 3
Known khat using history	Yes = 9
Evidence of using khat	Yes = 11; no = 3
Place of death	Home = 4; hospital = 6 (inc. 1 following traffic accident, and 1 in ambulance); railway station = 2; street = 1; outside flats = 1
Coroner's verdict	Self-harm = 2; suicide = 1; open = 2; accidental = 3; misadventure = 1; non-dependent abuse of drugs = 1; narrative = 2 (inc. 1 natural causes); abuse of drugs and natural causes = 1; unlawful killing = 1

UK deaths

Case	Role/association
1	Paranoid psychosis associated with history of khat use => traumatic suicide (none in body at death)
2	Possible suicide/accidental fall whilst judgement impaired (found in body)
3	Possible history of excessive use => traumatic murder and traumatic suicide
4	Paranoid schizophrenia exacerbated by khatting => accidental overdose
5	Ingestion of khat => high norephedrine levels => left ventricular failure => pulmonary oedema
6	Long term khat use => hepatic necrosis => sub-fulminant liver failure
7	Drug-induced psychosis/psychosis exacerbated by use of 'skunk' and khat over long period => traumatic suicide
8	Ingestion of khat (possibly no longer active), alcohol consumption => intoxication, impaired judgement/lack of co-ordination => traumatic road traffic accident (pedestrian)
9	Alcohol and khat in system => impaired judgement/co-ordination => loss of control of vehicle => traumatic road traffic accident (driver)
10	Overdose of injected heroin, but khat also in system
11	Abused khat => jaundice, night sweats, pyrexia => sub-acute liver failure
12	Ingestion of alcohol & khat +> intoxication +> fighting => collapse => cardiovascular event. Khat may have triggered an infarction or electrical instability (arrhythmia) causing death
13	Had consumed alcohol, cannabis & khat which may have contributed to his aggressive behaviour, leading to incident which triggered events leading to his assault and death
14	Excessive use of khat => fulminant hepatic necrosis => required liver transplant (failed)

UK deaths

The contribution of khat was varied. 50% traumatic in nature with external causes of death.

In 1 case, khat found in mouth, oesophagus & stomach but role unclear; decedent's khat-induced psychosis may have contributed to/or influenced fall from a height, or impaired judgement?

Impaired judgement due to use of khat & alcohol possibly instrumental in 2 cases where decedents fatally injured by contact with motor vehicles. Lack of motor co-ordination, impaired vision (Le Bras and Frétilière, 1965) or ability to judge speed properly (Khattab and Amer, 1995) may have contributed to the accidents.

UK deaths

Impaired judgement may have also impacted on case where the deceased was fatally assaulted as a result of being rude; had also consumed alcohol and cannabis.

‘Skunk’ use & khat contributed to problems experienced by one case. In another case khat present along with fatal levels of heroin.

Khat associated with psychiatric conditions in 4 cases (3 suicides + 1 murder), 1 accidental overdose. Earlier UK cases (Busby, 1987; Pantelis et al, 1989).

UK deaths

Physiological effects of khat consumption very clear in 5 cases. ?First reported fatalities due to khat toxicity since Heisch (1945).

Norephedrine/norpseudoephedrine producing adrenaline-like actions => increased likelihood of myocardial infarction => left ventricular failure and pulmonary oedema (Al-Motarreb et al. 1997, 2002, 2005; Croles et al., 2009; Health Canada, 2007).

Khat toxicity in 2 cases => hepatic necrosis and sub-fulminant liver failure; and in 3rd to sub-acute liver failure, in the presence of auto-immune hepatitis. These presentations consistent with findings for hepatitis (Brostoff et al., 2006; D'Souza et al., 2006) and liver disease (McCune et al., 2007). More recent papers confirm these findings and report additional deaths (Chapman et al., 2010; Peevers et al., 2010).

UK deaths

All male, aged 19-47 (mean = 36).

13/14 of East African descent: 12 Somali, 1 Eritrean.

However - one Polish: suggesting possibility of spread of khat use outside traditional ethnic khat-using populations.

All lived in areas with immigrant populations.

Length of residence in the UK 3-16 years (where known). 2 cases domiciled in UK because of the civil war in Somalia.

Case demographics are typical of khat users in the UK - relatively young, unemployed, Somali males, living with significant others. However, surprising no reported fatalities involving Yemenis or Ethiopians.

None were female – however

UK deaths

All male, aged 22-47 (mean = 36).

11/12 of East African descent: 10 Somali, 1 Eritrean.

However - one Polish: suggesting possibility of spread of khat use outside traditional ethnic khat-using populations.

All lived in areas with immigrant populations.

Length of residence in the UK 3-16 years (where known). 2 cases domiciled in UK because of the civil war in Somalia.

Case demographics are typical of khat users in the UK - relatively young, unemployed, Somali males, living with significant others. However, surprising no reported fatalities involving Yemenis.

Further cases

Another death, this time involving a Kuwaiti-born woman, presumably of Somali origin, who fell from a height from a flat in Northampton and died as a result, was reported by the media a few weeks ago. The death itself occurred in 2009. She had been chewing khat with a friend and her husband immediately before the incident.

I would be interested to hear details of any further cases so that we can follow them up with the relevant coroner. Any information provided will be treated with the utmost confidentiality, and will not be shared outside the team at St George's or the relevant coroner and his/her staff.

Discussion

Death can occur at any stage from the cultivation of khat to its consumption. Some of the factors can occur in combination, and it can be difficult sometimes to disentangle them.

Apart from poisoning from insecticide, traumatic deaths can be a feature of the cultivation, transportation, and trading activities undertaken by distributors and sellers. Violence can also occur in other settings.

Psychopathological effects – (i) impaired judgement leading to accidents and violence, (ii) causing or exacerbating psychoses or causing depression leading to suicide and even homicide. Physiological effects encompass (a) mechanical problems e.g. choking on pieces of the plant.; and (b) toxicity (i) causing heart problems leading to fatal heart attacks, and (ii) liver failure.

Discussion & conclusions

These UK cases illustrate some of the key issues related to the consumption of khat:

(a) psychological effects –

(i) impaired judgement leading to accidents and violence,

(ii) causing or exacerbating psychoses or causing depression leading to suicide and even homicide;

(b) physiological effects – toxicity is the primary concern here

(i) causing heart problems leading to fatal heart attacks, and

(ii) liver failure;

(c) mechanical problems e.g. choking on pieces of the plant; life-style aspects - part of the culture and thus being found in post-mortem toxicology.

Discussion & conclusions

This review of the UK experience and exploration of international research has demonstrated a lack of documented cases in the literature, as well as nothing by way of quantitative data.

These gaps in knowledge need to be filled.

This will lead to a much better understanding of the potential risks of death associated with khat use, based on empirical observation.

Need to identify and map sources of information on khat-related mortality, collate what is currently known in terms of statistics, and identify what gaps exist and how they might be filled.

An improved information-base enable the estimation of the possible numbers at risk of dying from the trade and use of khat.

Discussion & conclusions

Routine and systematic research of khat-using populations in respect of morbidity and mortality arising from khat is needed.

Only then can the best methods of supplying preventative and therapeutic interventions be considered in an informed way.

Meantime, dangers arising from khat use and its psychoactive constituents need to be brought to the attention of those in producing/growing countries, and those countries/regions that have become hosts to ever-increasing communities from these countries.

This will need to be specifically focused on target groups using a variety of media, including a variety of languages.

Discussion & conclusions

Lack of negative health results for khat in the literature should not lead to complacency or an assumption that khat use is free from toxic consequences. This absence of negative reports is due to a lack of population-based studies.

Whilst anecdotal reports are informative, systematic investigations using surveillance methodologies are needed to determine the incidence and prevalence of ill-effects of khat use.

The fatal case studies detailed above need to be borne in mind in future reviews of khat – but at the national and international levels - and the physiological and mental health risks associated with its use. These risks appear to be greater than previously realised. Something for policy-makers to chew on?

Comparison with other substances

This taxonomy provides a framework for describing cases and their different contexts.

These cases need to be related to the number and type(s) of khat users in order to assess risk(s) of morbidity and mortality.

This will facilitate the comparisons with the risks associated with other substances, particularly those which are chemically related such as methcathinones and amphetamine

Comparison with other substances

HUMAN PSYCHOPHARMACOLOGY
Hum. Psychopharmacol Clin Exp (2010)
Published online in Wiley InterScience
(www.interscience.wiley.com) DOI: 10.1002/hup.1090

An index of fatal toxicity for drugs of misuse

Leslie A. King* and John M. Corkery

27 Ivar Gardens, Basingstoke, RG24 8YD, UK

Objective To determine the lethal toxicity of five commonly-used illicit substances by relating the number of associated deaths to their availability.

Methods An index of toxicity was calculated for each of five drugs [heroin, cocaine/crack, ecstasy (MDMA), amphetamine and cannabis] as the ratio of the number of deaths associated with that substance to its availability in the period 2003–2007. Three separate proxy measures of availability were used (number of users as determined by household surveys, number of seizures by law enforcement agencies and estimates of the market size). All data are related to England and Wales only.

Results There was a broad correlation between all three denominators of availability. Not unexpectedly, heroin and cannabis showed, respectively, the highest and lowest toxicities. The index of fatal toxicity of MDMA was close to that of amphetamine and cocaine/crack. There was a rank correlation between this index and other measures of lethal toxicity based on safety ratios.

Conclusions These results are contrary to widely-held public views of the relative fatal toxicity of MDMA. Copyright © 2010 John Wiley & Sons, Ltd.

KEY WORDS—lethality; mortality statistics; drug users; law enforcement seizures; market size; illicit substances

Comparison with other substances

It might be possible to derive a lethality index value as calculated for other substances of mis(ue) by King and Corkery (2010). This was based on drug availability indicators, number of users, and number of fatalities.

An alternative approach might be to relate the number of fatalities where khat constituents are found in post mortem samples to the number of fatalities where these constituents and/or khat caused or contributed to death. This should be done for khat on its own and with other substances.

These measures would assist in informing the approach to assessing the need for regulation/control.

Acknowledgements

The National Programme on Substance Abuse Deaths would like to thank the individual coroners who kindly granted us access to the papers for the cases reported here. We are grateful to the coroners, Scottish Crime and Drug Enforcement Agency, and the General Register Office for Northern Ireland who regularly supply data to the National Programme on Substance Abuse Deaths. We are also grateful to the staff of the Office for National Statistics and the General Register Office for Scotland, as well as members of the London Toxicology Group, for checking their records. We are also grateful to colleagues in St George's, University of London, the European Monitoring Centre for Drugs and Drug Addiction in Lisbon, and the REITOX network for their assistance.

Funding

The surveillance work of the np-SAD is funded by the Department of Health (England)

References

See our original article for full references

Corkery, J.M., Schifano, F., Oyefeso, A., Ghodse, A.H., Tonia, T., Naidoo, V. and Button, J. (2010). ' 'Bundle of fun' or 'bunch of problems'? Case series of khat-related deaths in the UK'. *Drugs: education, prevention and policy*, Posted online on September 7, 2010. (doi:10.3109/09687637.2010.504200)

Contact details

John M Corkery, BA Hons (Open), MSc, MPhil
Senior Research Fellow in Drug Epidemiology
International Centre for Drug Policy
6th floor, Hunter Wing
St George's, University of London
Cranmer Terrace
London SW17 0RE

Tel: +44(0) 20 8725 2675
E-mail: jcorkery@sgul.ac.uk